

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
18 November 2004 (18.11.2004)

PCT

(10) International Publication Number
WO 2004/099195 A1

(51) International Patent Classification⁷: **C07D 409/12**,
417/12, 413/12, 411/12, 333/36, 333/28, A01N 43/10

(74) Agent: **WEICHSEL**, Christian; Syngenta Participations
AG, Intellectual Property, P.O. Box, CH-4002 Basel (CH).

(21) International Application Number:
PCT/EP2004/004194

(22) International Filing Date: 21 April 2004 (21.04.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0310464.3 7 May 2003 (07.05.2003) GB
0318920.6 12 August 2003 (12.08.2003) GB
0404806.2 3 March 2004 (03.03.2004) GB

(71) Applicant (for all designated States except US): **SYNGENTA PARTICIPATIONS AG** [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **EHRENFREUND, Josef** [AT/CH]; Syngenta Crop Protection AG, Schwarzwaldallee 215, CH-4058 Basel (CH). **WALTER, Harald** [DE/CH]; Syngenta Crop Protection AG, Schwarzwaldallee 215, CH-4058 Basel (CH). **TÖBLER, Hans** [CH/CH]; Syngenta Crop Protection AG, Schwarzwaldallee 215, CH-4058 Basel (CH). **LAMBERTH, Clemens** [DE/DE]; Syngenta Crop Protection, Schwarzwaldallee 215, CH-4058 Basel (CH).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HN, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SI, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: 3-CARBONYLAMINOTHIOPHENES AND THEIR USE AS FUNGICIDES



(57) Abstract: The invention also relates to novel compounds of formula (1) where Het is a 5- or 6-membered heterocyclic ring containing one to three heteroatoms, each independently selected from oxygen, nitrogen and sulphur, the ring being substituted by one to three groups R⁴; R¹ is hydrogen, optionally substituted (C₁₋₄)alkyl, formyl, optionally substituted (C₁₋₄)alkylC(=O), optionally substituted (C₁₋₄)alkylC(=O)O, optionally substituted (C₁₋₄)alkoxy(C₁₋₄)alkyl, optionally substituted

allyl, optionally substituted propargyl or optionally substituted allenyl; each R² is, independently, halogen, optionally substituted (C₁₋₄)alkyl, optionally substituted (C₁₋₄)alkoxy or optionally substituted (C₁₋₄)alkoxy(C₁₋₄)alkyl; R³ is either at position 2 or at position 4 of the thiophene ring and is an organic group containing three to thirteen carbon atoms and at least one silicon atom and, optionally, one to three heteroatoms, each independently selected from oxygen, nitrogen and sulphur, and is optionally substituted by one to four independently selected halogen atoms; each R⁴ is, independently, halogen, C₁₋₃ alkyl, C₁₋₃ haloalkyl, C₁₋₃alkoxy(C₁₋₃)alkyl or cyano; r is 0, 1 or 2; and X is O or S; or an N-oxide thereof, to novel intermediates used in the preparation of these compounds, to agrochemical compositions which comprise at least one of the novel compounds as an active ingredient and to the use of the active ingredients or compositions in agriculture or horticulture for controlling or preventing infestation of plants by phytopathogenic microorganisms, preferably fungi.


WO 2004/099195 A1

3-CARBONYLAMINOTHIOPHENES AND THEIR USE AS FUNGICIDES

The present invention relates to novel thiophene-3-amides, substituted in the 2- or 4-position of the thiophene ring by a silicon containing substituent, which have microbiocidal activity, in particular fungicidal activity. The invention also relates to the preparation of these compounds, to novel intermediates used in the preparation of these compounds, to agrochemical compositions which comprise at least one of the novel compounds as active ingredient, to the preparation of the compositions mentioned and to the use of the active ingredients or compositions in agriculture or horticulture for controlling or preventing infestation of plants by phytopathogenic microorganisms, preferably fungi.

Certain thiophene amides, substituted by a silicon containing substituent, are disclosed in US 5739338 and WO 97/19062.

The present invention provides a compound of formula (I)



where Het is a 5- or 6-membered heterocyclic ring containing one to three heteroatoms, each independently selected from oxygen, nitrogen and sulphur, the ring being substituted by one to three groups R^4 ; R^1 is hydrogen, optionally substituted (C_{1-4}) alkyl, formyl, optionally substituted (C_{1-4}) alkylC(=O), optionally substituted (C_{1-4}) alkylC(=O)O, optionally substituted (C_{1-4}) alkoxy (C_{1-4}) alkyl, optionally substituted allyl, optionally substituted propargyl or optionally substituted allenyl; each R^2 is, independently, halogen, optionally substituted (C_{1-4}) alkyl, optionally substituted (C_{1-4}) alkoxy or optionally substituted (C_{1-4}) alkoxy (C_{1-4}) alkyl;

R^3 is either at position 2 or at position 4 of the thiophene ring and is an organic group containing three to thirteen carbon atoms and at least one silicon atom and, optionally, one to three heteroatoms, each independently selected from oxygen, nitrogen and sulphur, and is optionally substituted by one to four independently selected halogen atoms; each R^4 is, independently, halogen, C_{1-3} alkyl, C_{1-3} haloalkyl, C_{1-3} alkoxy (C_{1-3}) alkyl or cyano; r is 0, 1 or 2; and X is O or S; or an N-oxide thereof.

In another aspect the present invention provides a compound of formula (I) as defined above provided that the Het ring is not 1,2,3-triazole when X is O.

Halogen is fluorine, chlorine, bromine or iodine, preferably fluorine, chlorine or bromine.

5 Each alkyl moiety is a straight or branched chain and is, for example, methyl, ethyl, *n*-propyl, *n*-butyl, *iso*-propyl, *n*-butyl, *sec*-butyl, *iso*-butyl or *tert*-butyl.

The alkenyl moieties, where appropriate, can be of either the (E)- or (Z)-configuration.

When present, each optional substituent on alkyl moieties, allyl, propargyl and allenyl is, independently, selected from halogen, hydroxy, cyano, carboxyl, methoxycarbonyl, 10 ethoxycarbonyl, methoxy, ethoxy, methylsulfonyl, ethylsulfonyl, difluoromethoxy, trifluoromethoxy and trifluorothiomethoxy.

Preferably R¹ is hydrogen, propargyl, allenyl, formyl, CH₃C(=O), C₂H₅C(=O) or CH₃OCH₂C(=O).

More preferably R¹ is hydrogen, propargyl, allenyl, CH₃C(=O), C₂H₅C(=O) or 15 CH₃OCH₂C(=O).

Most preferably R¹ is hydrogen.

Preferably, each R² is independently selected from halogen, methyl, trifluoromethyl and trifluoromethoxy.

It is preferred that Het is pyrazolyl, pyrrolyl, thiophenyl, furyl, thiazolyl, isothiazolyl, 20 oxazolyl, isoxazolyl, triazolyl, pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, 5.6-dihydropyranyl or 5.6-dihydro-1.4-oxathiinyl (more preferably pyrazolyl, pyrrolyl, thiophenyl, furyl, thiazolyl, oxazolyl, 1,2,3-triazolyl, pyridinyl, pyrimidinyl, pyridazinyl, 5.6-dihydropyranyl or 5.6-dihydro-1.4-oxathiinyl) each being substituted by one to three groups R⁴.

25 Preferably R³ is an aliphatic, saturated or unsaturated group containing three to thirteen carbon atoms and at least one silicon atom and, optionally, one to three heteroatoms, each independently selected from oxygen, nitrogen and sulphur, and is optionally substituted by one to four independently selected halogen atoms.

More preferably R³ is Y¹-Si(O_mMe)(O_nMe)(O_pY²) where m, n and p are each, 30 independently, 0 or 1; Y¹ is a bond or is alkandiyl (alkylene), alkendiyl (alkenylene) or alkindiyl (alkynylene), each of which is branched or unbranched and contains 1-6 carbon

atoms optionally interrupted by one or two oxygen atoms and optionally substituted by one to three independently selected halogen atoms; and Y^2 is alkyl or alkenyl, each of which is branched or unbranched and contains 1-5 carbon atoms optionally interrupted by one heteroatom selected from O, S and N and optionally substituted by one to three independently selected halogen atoms.

Even more preferably R^3 is $SiMe_3$, $SiMe_2Et$, $SiMe_2CHMe_2$, $SiMe_2CH_2CHMe_2$, $SiMe_2CH_2CMe_3$, $SiMe_2OCHMe_2$, $SiMe_2OCH_2CHMe_2$, CH_2SiMe_3 , CH_2SiMe_2Et , $CH_2SiMe_2CHMe_2$, $CH_2SiMe_2CH_2CHMe_2$, CH_2SiMe_2OMe , $CH_2SiMe_2OCHMe_2$, $CH_2SiMe_2OCH_2CHMe_2$, $CHMeSiMe_3$, $CHMeSiMe_2OMe$, $(CH_2)_2SiMe_3$, $(CH_2)_2SiMe_2Et$, $(CH_2)_2SiMe_2CHMe_2$, $(CH_2)_2SiMe_2CMe_3$, $(CH_2)_2SiMe_2CH_2CHMe_2$, $(CH_2)_2SiMe_2CH_2CH_2Me$, $(CH_2)_2SiMe_2CH_2CMe_3$, $(CH_2)_2SiMe_2OCHMe_2$, $(CH_2)_2SiMe_2OCH_2CHMe_2$, $CHMeCH_2SiMe_3$, $CHMeCH_2SiMe_2Et$, $CHMeCH_2SiMe_2CH_2CH_2Me$, $CHMeCH_2SiMe_2CHMe_2$, $CHMeCH_2SiMe_2CMe_3$, $CHMeCH_2SiMe_2CH_2CHMe_2$, $CFMeCH_2SiMe_3$, $CHMeCH_2CH_2SiMe_2OMe$, $CHMeCH_2SiMe_2OCHMe_2$, $CHMeCH_2SiMe_2OCH_2CHMe_2$, $CH_2CHMeSiMe_3$, $CH_2CHMeSiMe_2Et$, $CH_2CHMeSiMe_2CHMe_2$, $CHMeCHMeSiMe_3$, $CMe_2CH_2SiMe_3$, $(CH_2)_3SiMe_3$, $(CH_2)_3SiMe_2Et$, $(CH_2)_3SiMe_2CHMe_2$, $(CH_2)_3SiMe_2CH_2CHMe_2$, $(CH_2)_3SiMe_2OMe$, $(CH_2)_3SiMe_2OCHMe_2$, $(CH_2)_3SiMe_2OCH_2CHMe_2$, $CHMeCH_2CH_2SiMe_3$, $CHMeCH_2CH_2SiMe_2Et$, $CHMeCH_2CH_2SiMe_2CHMe_2$, $CHMeCH_2CH_2CH_2SiMe_2OMe$, $CHMeCH_2CH_2SiMe_2OCHMe_2$, $CMe=CHSiMe_3$ or $CH_2CH_2SiMe_2OMe$.

Most preferably R^3 is $SiMe_3$, $SiMe_2Et$, $SiMe_2CHMe_2$, $SiMe_2CH_2CHMe_2$, $SiMe_2CH_2CMe_3$, $(CH_2)_2SiMe_3$, $(CH_2)_2SiMe_2Et$, $(CH_2)_2SiMe_2CHMe_2$, $(CH_2)_2SiMe_2CMe_3$, $(CH_2)_2SiMe_2CH_2CHMe_2$, $CHMeCH_2SiMe_3$, $CHMeCH_2SiMe_2Et$, $CH_2CHMeSiMe_3$, $CMe_2CH_2SiMe_3$ or $(CH_2)_3SiMe_3$.

Nitrogen atoms in the Het ring are, independently, either unsubstituted or substituted by R^4 . When R^4 is a substituent on a nitrogen atom it is preferably C_{1-3} alkyl, C_{1-3} haloalkyl or methoxymethylene; more preferably C_{1-2} alkyl, CF_3 , CF_2Cl , CHF_2 , CH_2F or methoxymethylene; even more preferably methyl, CHF_2 or methoxymethylene; yet more preferably methyl or methoxymethylene; and most preferably methyl.

Carbon atoms in the Het ring which are not bonded to the atom substituted by $CXNR^1$ are, independently, either unsubstituted or substituted by R^4 . When R^4 is a substituent on a

carbon atom which is not bonded to the atom substituted by CXNR¹ it is preferably halogen, C₁₋₃ alkyl, C₁₋₃ haloalkyl or methoxymethylene; more preferably chloro, methoxymethylene, CH₃, CHF₂ or CF₃; yet more preferably chloro, CH₃, CHF₂ or CF₃; and even more preferably CH₃ or CF₃.

- 5 There may be one or two carbon atoms in the Het ring bonded to the atom substituted by CXNR¹; such carbon atoms are, independently, either unsubstituted or substituted by R⁴. When R⁴ is a substituent on a carbon atom bonded to the atom substituted by CXNR¹ it is preferably halogen, C₁₋₃ alkyl or C₁₋₃ haloalkyl; more preferably chloro, fluoro, bromo, C₁₋₂ alkyl, CF₃, CF₂Cl, CHF₂ or CH₂F; and even more preferably chloro, fluoro, bromo, methyl, CF₃, CHF₂ or CH₂F.

More preferably, when there is only one carbon atom in the Het ring bonded to the atom substituted by CXNR¹ that carbon atom is substituted by R⁴.

- More preferably, when there are two carbon atoms in the Het ring bonded to the atom substituted by CXNR¹ one such carbon atom is substituted by R⁴ and the other carbon atom is either unsubstituted or is substituted by fluoro, chloro or methyl.

Most preferably the combination of Het, R⁴ and CXNR¹ is 1-Me-3-R⁴-4-CXNR¹ pyrazolyl, 1-Me-3-R⁴-4-CXNR¹ pyrrolyl, 2-Me-3-R⁴-4-CXNR¹ thiazolyl or 2-R⁴-3-CXNR¹ pyridinyl.


Preferably X is oxygen.

- 20 Preferably r is 0.


When a compound of formula (I) is an N-oxide then it is preferred that Het is pyridinyl substituted by one to three groups R⁴.

Throughout this description, Me is used to represent the methyl group. Likewise, Et represents the ethyl group.

- 25 Thiophenes of formulae (IIa) and (IIb)



(IIa)



(IIb)

where R^3 is as defined above and A is NH_2 , $NHCH(O)$, optionally substituted $(C_{1-4})alkylC(=O)NH$, optionally substituted $(C_{1-4})alkylOC(=O)NH$, halogen or $N=C(C_6H_5)_2$ are useful as intermediates in the preparation of compounds of formula (I).

- 5 Compounds of formulae (IIa) and (IIb) where R^3 is $SiMe_2Et$, $SiMe_2CHMe_2$, $SiMe_2CH_2CHMe_2$, $SiMe_2CH_2CMe_3$, $CH=CHSiMe_3$, $(CH_2)_2SiMe_3$, $(CH_2)_2SiMe_2Et$, $(CH_2)_2SiMe_2CHMe_2$, $(CH_2)_2SiMe_2CMe_3$, $(CH_2)_2SiMe_2CH_2CHMe_2$, $CHMeCH_2SiMe_3$, $CHMeCH_2SiMe_2Et$, $CH_2CHMeSiMe_3$, $CMe_2CH_2SiMe_3$ or $(CH_2)_3SiMe_3$; and A is as defined above [provided that A is not halogen when R^3 is $SiMe_3$] are novel and are preferred as
10 intermediates for the preparation of compounds of formula (I). Compounds of formulae (IIa) and (IIb) where R^3 is $SiMe_3$ and A is halogen are already known.

- Therefore, in another aspect the present invention provides a compound of formula (IIa) or (IIb) where R^3 is $SiMe_3$, $SiMe_2Et$, $SiMe_2CHMe_2$, $SiMe_2CH_2CHMe_2$, $SiMe_2CH_2CMe_3$, $CH=CHSiMe_3$, $(CH_2)_2SiMe_3$, $(CH_2)_2SiMe_2Et$, $(CH_2)_2SiMe_2CHMe_2$, $(CH_2)_2SiMe_2CMe_3$,
15 $(CH_2)_2SiMe_2CH_2CHMe_2$, $CHMeCH_2SiMe_3$, $CHMeCH_2SiMe_2Et$, $CH_2CHMeSiMe_3$, $CMe_2CH_2SiMe_3$, $(CH_2)_3SiMe_3$; and A is NH_2 , $NHCH(O)$, optionally substituted $(C_{1-4})alkylC(=O)NH$, optionally substituted $(C_{1-4})alkylOC(=O)NH$, halogen or $N=C(C_6H_5)_2$; provided that A is not halogen when R^3 is $SiMe_3$.

- The compounds of formula (I), (IIa) and (IIb) may exist as different geometric or
20 optical isomers or in different tautomeric forms. This invention covers all such isomers and tautomers and mixtures thereof in all proportions as well as isotopic forms such as deuterated compounds.

- The compounds in Tables 1 to 28 below illustrate particularly preferred compounds of the invention, in which R^5 , R^6 and R^7 are each, independently, examples of R^4 as defined
25 above.

Table A represents Table 1 (when A is 1), represents Table 2 (when A is 2), represents Table 3 (when A is 3) and represents Table 4 (when A is 4).


Table A

Compound No.	R ¹	R ³	R ⁵	R ⁶	R ⁷	X
A.1	H	SiMe ₃	H	Me	CF ₃	O
A.2	H	SiMe ₃	H	Me	CF ₂ H	O
A.3	H	CH ₂ SiMe ₃	H	Me	CF ₃	O
A.4	H	CH ₂ SiMe ₃	H	Me	CF ₃	S
A.5	H	CH ₂ SiMe ₃	H	Me	CF ₂ H	O
A.6	propargyl	CH ₂ SiMe ₃	H	Me	CF ₃	O
A.7	H	CHMeSiMe ₃	H	Me	CF ₃	O
A.8	H	CHMeSiMe ₃	H	Me	CF ₂ H	O
A.9	H	CHMeSiMe ₃	H	Me	CF ₃	S
A.10	propargyl	CHMeSiMe ₃	H	Me	CF ₃	O
A.11	allenyl	CHMeSiMe ₃	H	Me	CF ₃	O
A.12	COMe	CHMeSiMe ₃	H	Me	CF ₃	O
A.13	H	CHMeSiMe ₃	F	Me	Me	O
A.14	H	(CH ₂) ₂ SiMe ₃	H	Me	CF ₃	O
A.15	H	(CH ₂) ₂ SiMe ₃	H	Me	CF ₃	S
A.16	H	(CH ₂) ₂ SiMe ₃	H	Me	CF ₂ H	O
A.17	propargyl	(CH ₂) ₂ SiMe ₃	H	Me	CF ₃	O
A.18	H	(CH ₂) ₂ SiMe ₃	F	Me	Me	O
A.19	H	(CH ₂) ₂ SiMe ₃	H	CH ₂ OMe	CF ₃	O
A.20	H	(CH ₂) ₂ SiMe ₃	H	CH ₂ OMe	CF ₂ H	O
A.21	H	CHMeCH ₂ SiMe ₃	H	Me	CF ₃	O
A.22	H	CHMeCH ₂ SiMe ₃	H	Me	CF ₃	S
A.23	H	CHMeCH ₂ SiMe ₃	H	CH ₂ OMe	CF ₃	O
A.24	H	CHMeCH ₂ SiMe ₃	H	Me	CF ₂ H	O
A.25	H	CHMeCH ₂ SiMe ₃	H	Me	CF ₂ H	S
A.26	propargyl	CHMeCH ₂ SiMe ₃	H	Me	CF ₃	O
A.27	allenyl	CHMeCH ₂ SiMe ₃	H	Me	CF ₃	O

A.28	propargyl	CHMeCH ₂ SiMe ₃	H	Me	CF ₂ H	O
A.29	allenyl	CHMeCH ₂ SiMe ₃	H	Me	CF ₂ H	O
A.30	H	CHMeCH ₂ SiMe ₃	F	Me	Me	O
A.31	COMe	CHMeCH ₂ SiMe ₃	H	Me	CF ₃	O
A.32	H	(CH ₂) ₃ SiMe ₃	H	Me	CF ₂ H	O
A.33	H	CH ₂ Si(Me ₂)Et	H	Me	CF ₃	O
A.34	H	CH ₂ Si(Me ₂)Et	H	Me	CF ₂ H	O
A.35	H	CH ₂ Si(Me ₂)CHMe ₂	H	Me	CF ₃	O
A.36	H	CH ₂ Si(Me ₂)CHMe ₂	H	Me	CF ₂ H	O
A.37	H	CH ₂ Si(Me ₂)OMe	H	Me	CF ₃	O
A.38	H	CH ₂ Si(Me ₂)OMe	H	Me	CF ₂ H	O
A.39	H	CH ₂ CH ₂ Si(Me ₂)OMe	H	Me	CF ₃	O
A.40	H	CHMeSi(Me ₂)OMe	H	Me	CF ₃	O
A.41	H	CHMeSi(Me ₂)OMe	H	Me	CF ₂ H	O
A.42	H	CH ₂ CH ₂ Si(Me ₂)OMe	H	Me	CF ₂ H	O
A.43	H	C(Me)=CHSiMe ₃	H	Me	CF ₃	O
A.44	H	SiMe ₃	H	Me	CH ₂ F	O
A.45	H	(CH ₂) ₂ SiMe ₃	H	Me	CH ₂ F	O
A.46	H	CH ₂ CHMeSiMe ₃	H	Me	CH ₂ F	O
A.47	H	CH ₂ CHMeSiMe ₃	H	Me	CF ₃	O
A.48	H	CH ₂ CHMeSiMe ₃	H	Me	CF ₂ H	O
A.49	H	CHMeCH ₂ SiMe ₃	H	Me	CH ₂ F	O
A.50	H	CMe ₂ CH ₂ SiMe ₃	H	Me	CF ₃	O
A.51	H	CMe ₂ CH ₂ SiMe ₃	H	Me	CF ₂ H	O
A.52	H	CHMeCHMeSiMe ₃	H	Me	CF ₂ H	O
A.53	H	CHMeCHMeSiMe ₃	H	Me	CF ₃	O
A.54	H	CH ₂ CMe ₂ SiMe ₃	H	Me	CF ₃	O
A.55	H	CH ₂ CMe ₂ SiMe ₃	H	Me	CF ₂ H	O
A.56	H	CHMe(CH ₂) ₂ SiMe ₃	H	Me	CF ₂ H	O
A.57	H	CHMe(CH ₂) ₂ SiMe ₃	H	Me	CF ₃	O

A.58	H	$(\text{CH}_2)_2\text{Si}(\text{Me}_2)(\text{CH}_2)_2\text{Me}$	H	Me	CF_2H	O
A.59	H	$(\text{CH}_2)_2\text{Si}(\text{Me}_2)(\text{CH}_2)_2\text{Me}$	H	Me	CF_3	O
A.60	H	$\text{CHMeCH}_2\text{Si}(\text{Me}_2)\text{CMe}_3$	H	Me	CF_3	O
A.61	H	$\text{C}(=\text{CH}_2)\text{CH}_2\text{Si}(\text{Me}_2)\text{CMe}_3$	H	Me	CF_3	O
A.62	H	$\text{C}(=\text{CH}_2)\text{CH}_2\text{Si}(\text{Me}_2)\text{CH}_2\text{Me}$	H	Me	CF_3	O
A.63	H	$(\text{CH}_2)_2\text{Si}(\text{Me}_2)\text{CH}_2\text{Me}$	H	Me	CF_3	O
A.64	H	$\text{CHMeCH}_2\text{Si}(\text{Me}_2)\text{CH}_2\text{Me}$	H	Me	CF_3	O
A.65	H	$(\text{CH}_2)_2\text{Si}(\text{Me}_2)\text{CHMe}_2$	H	Me	CF_3	O
A.66	H	$\text{CHMeCH}_2\text{Si}(\text{Me}_2)\text{CHMe}_2$	H	Me	CF_3	O
A.67	H	$\text{CHMeCH}_2\text{Si}(\text{Me}_2)\text{CH}_2\text{CHMe}_2$	H	Me	CF_3	O
A.68	H	$\text{Si}(\text{Me}_2)\text{CH}_2\text{Me}$	H	Me	CF_2H	O
A.69	H	$\text{Si}(\text{Me}_2)\text{CH}_2\text{Me}$	H	Me	CF_3	O
A.70	H	$\text{Si}(\text{Me}_2)\text{CHMe}_2$	H	Me	CF_3	O
A.71	H	$\text{Si}(\text{Me}_2)\text{CHMe}_2$	H	Me	CF_2H	O
A.72	H	$\text{Si}(\text{Me}_2)\text{CH}_2\text{CHMe}_2$	H	Me	CF_2H	O
A.73	H	$\text{Si}(\text{Me}_2)\text{CH}_2\text{CHMe}_2$	H	Me	CF_3	O
A.74	H	$\text{C}:\text{CCH}_2\text{SiMe}_3$	H	Me	CF_3	O
A.75	propargyl	$(\text{CH}_2)_2\text{SiMe}_3$	H	Me	CF_2H	O
A.76	allenyl	$(\text{CH}_2)_2\text{SiMe}_3$	H	Me	CF_2H	O
A.77	allenyl	$(\text{CH}_2)_2\text{SiMe}_3$	H	Me	CF_3	O
A.78	H	$(\text{CH}_2)_2\text{SiMe}_3$	H	Me	CF_2Cl	O
A.79	H	$(\text{CH}_2)_3\text{SiMe}_3$	H	Me	CF_3	O
A.80	H	$(\text{CH}_2)_2\text{SiMe}_3$	Br	Me	CF_3	O
A.81	H	$(\text{CH}_2)_2\text{SiMe}_3$	Cl	Me	CF_3	O
A.82	H	$(\text{CH}_2)_2\text{SiMe}_3$	H	Me	Me	O
A.83	H	$\text{C}:\text{CSiMe}_3$	H	Me	CF_3	O
A.84	H	$\text{C}:\text{CSiMe}_3$	H	Me	CF_2H	O
A.85	CHO	$(\text{CH}_2)_2\text{SiMe}_3$	H	Me	CF_2H	O
A.86	CHO	$(\text{CH}_2)_2\text{SiMe}_3$	H	Me	CF_3	O

Table 1 provides 86 compounds of formula (Ia) where R^1 , R^3 , R^5 , R^6 , R^7 and X are as defined in Table 1.



5

Table 2 provides 86 compounds of formula (Ib) where R^1 , R^3 , R^5 , R^6 , R^7 and X are as defined in Table 2.




Table 3 provides 86 compounds of formula (Ic) where R^1 , R^3 , R^5 , R^6 , R^7 and X are as defined in Table 3.





Table 4 provides 86 compounds of formula (Id) where R^1 , R^3 , R^5 , R^6 , R^7 and X are as defined in Table 4.



15


Table B represents Table 5 (when B is 5), represents Table 6 (when B is 6), represents Table 7 (when B is 7) and represents Table 8 (when B is 8).

Table B

Compound No.	R ¹	R ³	R ⁵	R ⁶	X
B.1	H	SiMe ₃	Me	CF ₃	O
B.2	H	SiMe ₃	Me	CF ₂ H	O
B.3	H	CH ₂ SiMe ₃	Me	CF ₃	O
B.4	H	CH ₂ SiMe ₃	Me	CF ₃	S
B.5	H	CH ₂ SiMe ₃	Me	CF ₂ H	O
B.6	propargyl	CH ₂ SiMe ₃	Me	CF ₃	O
B.7	H	CHMeSiMe ₃	Me	CF ₃	O
B.8	H	CHMeSiMe ₃	Me	CF ₂ H	O
B.9	H	CHMeSiMe ₃	Me	CF ₃	S
B.10	propargyl	CHMeSiMe ₃	Me	CF ₃	O
B.11	allenyl	CHMeSiMe ₃	Me	CF ₃	O
B.12	COMe	CHMeSiMe ₃	Me	CF ₃	O
B.13	H	CHMeSiMe ₃	Me	Me	O
B.14	H	(CH ₂) ₂ SiMe ₃	Me	CF ₃	O
B.15	H	(CH ₂) ₂ SiMe ₃	Me	CF ₃	S
B.16	H	(CH ₂) ₂ SiMe ₃	Me	CF ₂ H	O
B.17	propargyl	(CH ₂) ₂ SiMe ₃	Me	CF ₃	O
B.18	H	(CH ₂) ₂ SiMe ₃	Me	Me	O
B.19	H	(CH ₂) ₂ SiMe ₃	CF ₃	CF ₃	O
B.20	H	CHMeCH ₂ SiMe ₃	Me	CF ₃	O
B.21	H	CHMeCH ₂ SiMe ₃	Me	CF ₃	S
B.22	H	CHMeCH ₂ SiMe ₃	Me	CF ₂ H	O
B.23	H	CHMeCH ₂ SiMe ₃	Me	CF ₂ H	S
B.24	propargyl	CHMeCH ₂ SiMe ₃	Me	CF ₃	O
B.25	propargyl	CHMeCH ₂ SiMe ₃	Me	CF ₂ H	O

B.26	H	CHMeCH ₂ SiMe ₃	Me	Me	O
B.27	H	CHMeCH ₂ SiMe ₃	CF ₃	CF ₃	O
B.28	COMe	CHMeCH ₂ SiMe ₃	Me	CF ₃	O
B.29	H	(CH ₂) ₃ SiMe ₃	Me	CF ₃	O
B.30	H	(CH ₂) ₃ SiMe ₃	Me	CF ₂ H	O
B.31	H	CH ₂ Si(Me ₂)Et	Me	CF ₃	O
B.32	H	CH ₂ Si(Me ₂)Et	Me	CF ₂ H	O
B.33	H	CH ₂ Si(Me ₂)CHMe ₂	Me	CF ₃	O
B.34	H	CH ₂ Si(Me ₂)CHMe ₂	Me	CF ₂ H	O
B.35	H	CH ₂ CHMeSiMe ₃	Me	CF ₃	O
B.36	H	CH ₂ CHMeSiMe ₃	Me	CF ₂ H	O
B.37	H	CMe ₂ CH ₂ SiMe ₃	Me	CF ₃	O
B.38	H	CMe ₂ CH ₂ SiMe ₃	Me	CF ₂ H	O
B.39	H	CHMeCHMeSiMe ₃	Me	CF ₂ H	O
B.40	H	CHMeCHMeSiMe ₃	Me	CF ₃	O
B.41	H	CH ₂ CMe ₂ SiMe ₃	Me	CF ₃	O
B.42	H	CH ₂ CMe ₂ SiMe ₃	Me	CF ₂ H	O
B.43	H	CHMe(CH ₂) ₂ SiMe ₃	Me	CF ₂ H	O
B.44	H	CHMe(CH ₂) ₂ SiMe ₃	Me	CF ₃	O
B.45	H	(CH ₂) ₂ SiMe ₃	CH ₂ OMe	CH ₂ Me	O
B.46	H	(CH ₂) ₂ SiMe ₃	CH ₂ OCH ₂ Me	CH ₂ Me	O
B.47	H	SiMe ₂ CH ₂ CHMe ₂	Me	CF ₃	O


Table 5 provides 47 compounds of formula (Ie) where R¹, R³, R⁵, R⁶ and X are as defined in Table 5.




(Ie)

- 12 -

Table 6 provides 47 compounds of formula (If) where R^1 , R^3 , R^5 , R^6 and X are as defined in Table 6.



5 Table 7 provides 47 compounds of formula (Ig) where R^1 , R^3 , R^5 , R^6 and X are as defined in Table 7.



10 Table 8 provides 47 compounds of formula (Ih) where R^1 , R^3 , R^5 , R^6 and X are as defined in Table 8.





Table BB represents Table 9 (when BB is 9) and represents Table 10 (when BB is 10).

Table BB


Compound No.	R ¹	R ³	R ⁵	R ⁶	X
BB.1	H	CH ₂ SiMe ₃	Me	CF ₃	S
BB.2	H	CHMeSiMe ₃	Me	CF ₃	S
BB.3	H	(CH ₂) ₂ SiMe ₃	Me	CF ₃	S
BB.4	H	CHMeCH ₂ SiMe ₃	Me	CF ₃	S
BB.5	H	CHMeCH ₂ SiMe ₃	Me	CF ₂ H	S

Table 9 provides 5 compounds of formula (Ii) where R¹, R³, R⁵, R⁶ and X are as defined in Table 9.



5

Table 10 provides 5 compounds of formula (Ij) where R¹, R³, R⁵, R⁶ and X are as defined in Table 10.



10


Table C represents Table 11 (when C is 11), represents Table 12 (when C is 12), represents Table 13 (when C is 13) and represents Table 14 (when C is 14).

Table C

Compound No.	R ¹	R ³	R ⁵	R ⁶	R ⁷	X
C.1	H	SiMe ₃	Me	Me	H	O
C.2	H	SiMe ₃	Me	Me	H	O
C.3	H	CH ₂ SiMe ₃	Me	Me	Me	O
C.4	H	CH ₂ SiMe ₃	Me	Me	CF ₃	O
C.5	H	CH ₂ SiMe ₃	Me	Me	H	O
C.6	propargyl	CH ₂ SiMe ₃	Me	Me	CF ₃	O
C.7	H	CHMeSiMe ₃	Me	Me	CF ₃	O
C.8	H	CHMeSiMe ₃	Me	Me	Me	O
C.9	H	CHMeSiMe ₃	Me	Me	Me	S
C.10	propargyl	CHMeSiMe ₃	Me	Me	Me	O
C.11	allenyl	CHMeSiMe ₃	Me	Me	Me	O
C.12	COMe	CHMeSiMe ₃	Me	Me	Me	O
C.13	H	CHMeSiMe ₃	Me	Me	Me	O
C.14	H	(CH ₂) ₂ SiMe ₃	Me	Me	CF ₃	O
C.15	H	(CH ₂) ₂ SiMe ₃	H	H	CF ₃	O
C.16	H	(CH ₂) ₂ SiMe ₃	H	H	CF ₃	S
C.17	propargyl	(CH ₂) ₂ SiMe ₃	H	H	CF ₃	O
C.18	H	(CH ₂) ₂ SiMe ₃	Me	Me	H	O
C.19	H	CHMeCH ₂ SiMe ₃	H	H	CF ₃	O
C.20	H	CHMeCH ₂ SiMe ₃	H	H	CF ₃	S
C.21	H	CHMeCH ₂ SiMe ₃	Me	Me	Me	O
C.22	H	CHMeCH ₂ SiMe ₃	H	Me	CF ₃	O
C.23	H	CHMeCH ₂ SiMe ₃	Me	Me	H	O
C.24	COMe	CHMeCH ₂ SiMe ₃	Me	Me	H	O
C.25	propargyl	CHMeCH ₂ SiMe ₃	Me	Me	H	O
C.26	allenyl	CHMeCH ₂ SiMe ₃	Me	Me	H	O
C.27	propargyl	CHMeCH ₂ SiMe ₃	Me	Me	Me	O

C.28	allenyl	CHMeCH ₂ SiMe ₃	Me	Me	Me	O
C.29	COMe	CHMeCH ₂ SiMe ₃	Me	Me	Me	O
C.30	COEt	CHMeCH ₂ SiMe ₃	Me	Me	Me	O
C.31	H	CH ₂ CHMeSiMe ₃	H	H	CF ₃	O
C.32	H	CH ₂ CHMeSiMe ₃	H	H	CF ₃	S
C.33	H	CH ₂ CHMeSiMe ₃	Me	Me	Me	O
C.34	H	CH ₂ CHMeSiMe ₃	H	Me	CF ₃	O
C.35	H	CH ₂ CHMeSiMe ₃	Me	Me	H	O

Table 11 provides 35 compounds of formula (Ik) where R¹, R³, R⁵, R⁶, R⁷ and X are as defined in Table 11.



5 Table 12 provides 35 compounds of formula (II) where R¹, R³, R⁵, R⁶, R⁷ and X are as defined in Table 12.




Table 13 provides 35 compounds of formula (Im) where R¹, R³, R⁵, R⁶, R⁷ and X are as defined in Table 13.





Table 14 provides 35 compounds of formula (In) where R^1 , R^3 , R^5 , R^6 , R^7 and X are as defined in Table 14.




- 5 Table D represents Table 15 (when D is 15), represents Table 16 (when D is 16), represents Table 17 (when D is 17) and represents Table 18 (when D is 18).

Table D

Compound No.	R^1	R^3	R^5	X
D.1	H	SiMe ₃	Me	O
D.2	H	SiMe ₃	CF ₃	O
D.3	H	CH ₂ SiMe ₃	Me	O
D.4	H	CH ₂ SiMe ₃	CF ₃	S
D.5	COMe	CH ₂ SiMe ₃	Me	O
D.6	propargyl	CH ₂ SiMe ₃	Me	O
D.7	H	CHMeSiMe ₃	Me	O
D.8	H	CHMeSiMe ₃	CF ₃	O
D.9	H	CHMeSiMe ₃	CF ₃	S
D.10	propargyl	CHMeSiMe ₃	Me	O
D.11	allenyl	CHMeSiMe ₃	Me	O
D.12	COMe	CHMeSiMe ₃	Me	O
D.13	propargyl	CHMeSiMe ₃	CF ₃	O
D.14	H	(CH ₂) ₂ SiMe ₃	Me	O
D.15	H	(CH ₂) ₂ SiMe ₃	CF ₃	O
D.16	H	(CH ₂) ₂ SiMe ₃	CF ₃	S
D.17	propargyl	(CH ₂) ₂ SiMe ₃	Me	O

D.18	COMe	$(\text{CH}_2)_2\text{SiMe}_3$	Me	O
D.19	H	$\text{CHMeCH}_2\text{SiMe}_3$	Me	O
D.20	H	$\text{CHMeCH}_2\text{SiMe}_3$	CF_3	O
D.21	H	$\text{CHMeCH}_2\text{SiMe}_3$	CF_3	S
D.22	propargyl	$\text{CHMeCH}_2\text{SiMe}_3$	Me	O
D.23	allenyl	$\text{CHMeCH}_2\text{SiMe}_3$	Me	O
D.24	COMe	$\text{CHMeCH}_2\text{SiMe}_3$	Me	O
D.25	propargyl	$\text{CHMeCH}_2\text{SiMe}_3$	CF_3	O
D.26	allenyl	$\text{CHMeCH}_2\text{SiMe}_3$	CF_3	O
D.27	COMe	$\text{CHMeCH}_2\text{SiMe}_3$	CF_3	O
D.28	allenyl	$\text{CHMeCH}_2\text{SiMe}_3$	Me	O
D.29	H	$(\text{CH}_2)_3\text{SiMe}_3$	Me	O
D.30	H	$(\text{CH}_2)_3\text{SiMe}_3$	CF_3	O
D.31	H	$\text{CH}_2\text{CHMeSiMe}_3$	Me	O
D.32	H	$\text{CH}_2\text{CHMeSiMe}_3$	CF_3	O

Table 15 provides 32 compounds of formula (Io) where R^1 , R^3 , R^5 and X are as defined in Table 15.



5 Table 16 provides 32 compounds of formula (Ip) where R^1 , R^3 , R^5 and X are as defined in Table 16.




Table 17 provides 32 compounds of formula (Iq) where R^1 , R^3 , R^5 and X are as defined in Table 17.




Table 18 provides 32 compounds of formula (Ir) where R^1 , R^3 , R^5 and X are as defined in Table 18.




Table E represents Table 19 (when E is 19), represents Table 20 (when E is 20), represents Table 21 (when E is 21), represents Table 22 (when E is 22), represents Table 23 (when E is 23) and represents Table 24 (when E is 24).


Table E

Compound No.	R^1	R^3	R^5	X
E.1	H	SiMe ₃	Cl	O
E.2	H	SiMe ₃	CF ₃	O
E.3	H	CH ₂ SiMe ₃	Cl	O
E.4	H	CH ₂ SiMe ₃	Br	O
E.5	H	CH ₂ SiMe ₃	CF ₃	O
E.6	propargyl	CH ₂ SiMe ₃	Cl	O
E.7	H	CHMeSiMe ₃	Cl	O

E.8	H	CHMeSiMe ₃	Br	O
E.9	H	CHMeSiMe ₃	CF ₃	O
E.10	propargyl	CHMeSiMe ₃	Cl	O
E.11	allenyl	CHMeSiMe ₃	Cl	O
E.12	COMe	CHMeSiMe ₃	Cl	O
E.13	H	CHMeSiMe ₃	Cl	S
E.14	H	(CH ₂) ₂ SiMe ₃	Cl	O
E.15	H	(CH ₂) ₂ SiMe ₃	Br	O
E.16	H	(CH ₂) ₂ SiMe ₃	CF ₃	O
E.17	propargyl	(CH ₂) ₂ SiMe ₃	Cl	O
E.18	COMe	(CH ₂) ₂ SiMe ₃	Cl	O
E.19	H	CHMeCH ₂ SiMe ₃	Cl	O
E.20	H	CHMeCH ₂ SiMe ₃	Cl	S
E.21	H	CHMeCH ₂ SiMe ₃	Br	O
E.22	H	CHMeCH ₂ SiMe ₃	CF ₃	O
E.23	propargyl	CHMeCH ₂ SiMe ₃	Cl	O
E.24	allenyl	CHMeCH ₂ SiMe ₃	Cl	O
E.25	COMe	CHMeCH ₂ SiMe ₃	Cl	O
E.26	propargyl	CHMeCH ₂ SiMe ₃	Br	O
E.27	allenyl	CHMeCH ₂ SiMe ₃	Br	O
E.28	COMe	CHMeCH ₂ SiMe ₃	Br	O
E.29	COCH ₂ OMe	CHMeCH ₂ SiMe ₃	Cl	O
E.30	COCH ₂ OMe	CHMeCH ₂ SiMe ₃	CF ₃	O
E.31	H	(CH ₂) ₃ SiMe ₃	Cl	O
E.32	H	(CH ₂) ₃ SiMe ₃	Br	O
E.33	H	(CH ₂) ₃ SiMe ₃	CF ₃	O
E.34	H	CH ₂ CHMeSiMe ₃	CF ₃	O
E.35	H	CH ₂ CHMeSiMe ₃	Cl	O
E.36	H	CH ₂ CHMeSiMe ₃	Br	O
E.37	H	SiMe ₂ CH ₂ Me	CF ₃	O

E.38	H	SiMe ₂ CH ₂ Me	Cl	O
E.39	H	SiMe ₂ CH ₂ Me	Br	O
E.40	H	SiMe ₂ CHMe ₂	CF ₃	O
E.41	H	SiMe ₂ CHMe ₂	Cl	O
E.42	H	SiMe ₂ CHMe ₂	Br	O
E.43	H	SiMe ₂ CH ₂ CH ₂ Me	CF ₃	O
E.44	H	SiMe ₂ CH ₂ CH ₂ Me	Cl	O
E.45	H	SiMe ₂ CH ₂ CH ₂ Me	Br	O

Table 19 provides 45 compounds of formula (Is) where R¹, R³, R⁵ and X are as defined in Table 19.



5 Table 20 provides 45 compounds of formula (It) where R¹, R³, R⁵ and X are as defined in Table 20.




Table 21 provides 45 compounds of formula (Iu) where R¹, R³, R⁵ and X are as defined in Table 21.




Table 22 provides 45 compounds of formula (Iv) where R^1 , R^3 , R^5 and X are as defined in Table 22.




Table 23 provides 45 compounds of formula (Iw) where R^1 , R^3 , R^5 and X are as defined in Table 23.




Table 24 provides 45 compounds of formula (Ix) where R^1 , R^3 , R^5 and X are as defined in Table 24.





Table 25 provides 10 compounds of formula (Iy) where R^7 and $(R^2)_r$ are as defined in Table 25.




- 22 -

Table 25

Compound Number	R ⁷	(R ²) _r
25.1	CF ₃	2-Cl
25.2	CF ₃	5-Cl
25.3	CF ₃	2-Br
25.4	CF ₃	5-Br
25.5	CF ₂ H	2-Cl
25.6	CF ₂ H	5-Cl
25.7	CF ₂ H	2-Br
25.8	CF ₂ H	5-Br
25.9	CF ₃	2-Cl,5-Cl
25.10	CF ₂ H	2-Cl,5-Cl

Table 26 provides 10 compounds of formula (Iz) where R⁷ and (R²)_r are as defined in Table 26.



(Iz)

5

Table 26

Compound Number	R ⁷	(R ²) _r
26.1	CF ₃	4-Cl
26.2	CF ₃	5-Cl
26.3	CF ₃	4-Br
26.4	CF ₃	5-Br

- 23 -


26.5	CF ₃	4-Cl,5-Cl
26.6	CF ₂ H	4-Cl
26.7	CF ₂ H	5-Cl
26.8	CF ₂ H	4-Br
26.9	CF ₂ H	5-Br
26.10	CF ₂ H	4-Cl,5-Cl

Table G represents Table 27 (when G is 27) and represents Table 28 (when G is 28).

Table G


Compound Number	R ³	A
G.1	(CH ₂) ₂ SiMe ₃	NH ₂
G.2	(CH ₂) ₂ SiMe ₃	NHCHO
G.3	(CH ₂) ₂ SiMe ₃	NHCOOC(CH ₃) ₃
G.4	(CH ₂) ₂ SiMe ₃	Br
G.5	(CH ₂) ₂ SiMe ₃	N=C(C ₆ H ₅) ₂
G.6	(CH ₂) ₂ SiMe ₃	NHCOC(CH ₃) ₃
G.7	CHMeCH ₂ SiMe ₃	NH ₂
G.8	CHMeCH ₂ SiMe ₃	NHCHO
G.9	CHMeCH ₂ SiMe ₃	NHCOOC(CH ₃) ₃
G.10	CHMeCH ₂ SiMe ₃	Br
G.11	CHMeCH ₂ SiMe ₃	N=C(C ₆ H ₅) ₂
G.12	CHMeCH ₂ SiMe ₃	NHCOC(CH ₃) ₃
G.13	SiMe ₃	NHCHO
G.14	SiMe ₂ CH ₂ CHMe ₂	NHCHO

Table 27 provides 14 compounds of formula (IIa) where R³ and A are as defined in Table 27.



(IIa)

5 Table 28. Table 28 provides 14 compounds of formula (IIb) where R³ and A are as defined in



(IIb)

Throughout this description, temperatures are given in degrees Celsius; "NMR" means nuclear magnetic resonance spectrum; MS stands for mass spectrum; and "%" is percent by weight, unless corresponding concentrations are indicated in other units.

10 The following abbreviations are used throughout this description:

m.p. = melting point	b.p.= boiling point.
s = singlet	br = broad
d = doublet	dd = doublet of doublets
t = triplet	q = quartet
m = multiplet	ppm = parts per million
qd = quartet of doublets	sext = sextet

Table 29 shows selected melting point, selected NMR data, all with CDCl₃ as the solvent (unless otherwise stated; if a mixture of solvents is present, this is indicated as, for example, (CDCl₃ / d₆-DMSO)) and characteristic mass spectrum signals (no attempt is made to list all characterising data in all cases) for compounds of Tables 1 to 28.

15

Table 29

Compound Number	¹ H-NMR data: (ppm/multiplicity/number of Hs) or mass spectrum signal	m.p. / (°C)
1.14		140-142
1.16		145-147
2.1		121-124
2.2		85-88
2.14		109-111
2.16		140-142
2.72		57-59
2.73		70-71
2.75		80-83
2.76		74-76
2.84		130-133
2.85	(M ⁺ + Cl) : 420	
6.1		88-92
6.14		74-77
6.47	(M ⁺ + H) : 407	
20.1	0.4 (s,9); 7.45 (m,1); 7.6 (d,1); 7.8 (d,1); 8.2 (br,1); 8.25 (m,1); 8.6 (m,1).	
20.14		106-110
27.2	(M ⁺ -1) : 226	
27.4	0.0(s,9); 0.8(m,2); 2.6(m,2); 6.9(d,1); 7.2(d,1)	126-128
28.1	0.05(s,9); 0.85(m,2); 2.6(m,2); 3.3(br.s,2); 6.6(d,1); 6.9(d,1)	
28.2		52-53
28.3	0.0(s,9); 0.9(m,2); 1.4(s,9); 2.6(m,2); 6.0(br.s,1); 7.0(m,2)	

The compounds according to the present invention may be prepared according to the following reaction schemes, in which, unless otherwise stated, the definition of each variable is as defined above for a compound of formula (I).

There are a number of alternative methods for preparing a compound of formula (I).

Method A

A compound of formula (I) may be prepared by reacting a compound of formula (IIa) or (IIb) [in which A is NH₂, NHCH(O), optionally substituted (C₁₋₄)alkylC(=O)NH or optionally substituted (C₁₋₄)alkylOC(=O)NH] with a compound of formula Het-C(=O)OR' [where R' is C₁₋₅ alkyl] in the presence of a strong base [for example NaH or sodium hexamethyldisilazane], in a dry polar solvent [preferably THF] and at a temperature between -10°C and the boiling point of the solvent [preferably at ambient temperature]. The article by

J.Wang et al [Synlett 2001, 1485] provides details of analogous preparations. When A is NHCH(O), optionally substituted (C₁₋₄)alkylC(=O)NH or optionally substituted (C₁₋₄)alkylOC(=O)NH; and a compound of formula (I) in which R¹ is H is desired, then hydrolysis according to Method E below must follow.

5 Method B

A compound of formula (I) may be prepared by reacting a compound of formula (IIa) or (IIb) [where A is as defined above in Method A] with a compound of formula Het-C(=O)R'' [where R'' is OH or a leaving group, such as Cl, Br, F or OC(=O)C₁₋₄ alkyl] in an inert organic solvent [such as ethylacetate, dichloromethane, dioxane, THF or DMF] and at a
10 temperature between -10°C and the boiling point of the solvent [preferably at ambient temperature]. If R'' is OH, then the reaction is carried out in the presence of an activating agent [for example BOP-Cl] and two equivalents of a base [such as a tertiary amine, an inorganic carbonate or a hydrogen carbonate]. Alternatively, if R'' is a leaving group, then the reaction is carried out in the presence of at least one equivalent of base [for example pyridine,
15 a tertiary amine, an inorganic carbonate or a hydrogen carbonate; a stronger base, such as NaH or sodium hexamethyldisilazane is used when A is NHCH(O), optionally substituted (C₁₋₄)alkylC(=O)NH or optionally substituted (C₁₋₄)alkylOC(=O)NH]. If a compound of formula (I) in which R¹ is H is desired, then hydrolysis according to Method E must follow.

Method C

20 A compound of formula (I) [where R¹ is as defined above but is not hydrogen] may be prepared by reacting a compound of formula (I) [where R¹ is hydrogen] with a compound of formula R¹-L¹ [where R¹ is as defined above but is not hydrogen; and L¹ is a leaving group, such as Cl, Br, I, a sulfonate (for example a mesylate or a tosylate) or OC(O)C₁₋₄ alkyl] in a solvent [such as an halogenated solvent (for example dichloromethane), an ether, ethylacetate,
25 DMF or even water (as a biphasic mixture, optionally in the presence of a phase transfer catalyst such as tetrabutylammonium hydrogensulfate)] and in the presence of a base [such as a tertiary amine, an alkali carbonate, an alkali bicarbonate, an alkali hydroxide or NaH; though when L¹ is O(CO)C₁₋₄ alkyl then simply heating without base is possible].

Method D

30 A compound of formula (I) may be prepared by reacting a compound of formula (IIa) or (IIb) [where A is halogen, preferably bromo or iodo] with a compound of formula

Het-C(=O)NH₂ in the presence of a Cu(I) compound and an aprotic solvent [such as a cyclic ether, for example dioxane] at an elevated temperature and preferably at reflux. It is preferred that CuI is used at 2% to 100% mole/mole, relative to the compound of formula (IIb), in the presence of a 1,2-diamine (such as 1,2-diamino cyclohexane or ethylene diamine) as a ligand-
 5 forming substance and at least one equivalent of a base (such as an alkali carbonate or an alkali phosphate). The article by A.Klapars et al. J.Am.Chem.Soc. 123, 7727 (2001) provides details of analogous preparations.


Method E

A compound of formula (I) [where R¹ is H] may be prepared from a compound of
 10 formula (I) [where R¹ is as defined above, but is not H] by acidic or alkaline hydrolysis. For this purpose the compound is treated with aqueous acid or base, for example HCl, HBr or an organic hydroxide [such as sodium-, potassium-, calcium- or barium-hydroxide] in an appropriate solvent which is preferably mixable with water [for example THF, dioxane, a lower alcohol or water itself] at ambient or elevated temperatures.

15 Method F

A compound of formula (I) [where R¹ is H] may be prepared from a compound of
 formula (IIa) or (IIb) [where A is N=C(C₆H₅)₂] by converting A to NH₂, for example according to methods described by J. Ahman et al. Tetrahedron Letters 38, 6363 (1997) and proceeding according to Method A or Method B, preferably without isolation or purification
 20 of the intermediates:

Some compounds of formulae (IIa) and (IIb) are already known; novel compounds of formula (IIa) or (IIb) may be prepared according to the following synthetic strategies which are depicted in the following scheme and described below:



Step 1: Starting from a suitable precursor of formula (III) which carries a precursor substituent PS [which is a protected or free amino function, a substituent which may be converted to NH₂ in a later stage of the synthesis or a precursor group for A as defined above] and, optionally, a substituent T [which is convertible to "Si", a silicon containing substituent which is either R³ as defined above or is a precursor for such R³] an appropriate Si-containing functionality ("Si") is introduced into the ortho-position to give a compound of formula (IV).

Step 2: If necessary, the introduced "Si" group is further manipulated to form the desired substituent R³ in a compound of formula (V).

Step 3: Deprotection if necessary or conversion of the precursor substituent to yield substituent A, giving a compound of formula (IIa) or (IIb).

Steps 2 and 3 may also be carried out in reversed order.

Examples of protecting groups for the NH₂ functionality are formyl, acyl, haloacyl, trialkylsilyl, (substituted)benzyl and alkoxycarbonyl. A more comprehensive list of methods for protection and deprotection of aromatic and heteroaromatic amines which are useful in the context of the present invention may be found in T.W. Green and P.G.M. Wuts, Protective Groups in Organic Synthesis 3rd edition p.503-614 (Wiley 1999).

Examples for precursor substituents PS are nitro and azido [both of which may be converted to NH₂ by reduction or hydrogenation], carboxyl and carboxy derivatives [which may undergo rearrangements to form isocyanates, for example by Curtius-, Schmidt- or Hofmann-degradation] and halides and triflates [which may be converted to NH₂ in protected or unprotected form via palladium catalysed amination reactions currently known under the name "Buchwald Hartwig" reaction, see for example J. Ahman et al. Tetrahedron Letters 38,6363 (1997) and X.Huang et al., Org.Lett. 3, 3417 (2001) and references cited therein].

More comprehensive lists for useful precursor substituents for NH₂ may be found in Rodd's Chemistry of Carbon Compounds III B and its supplements (Elsevier 1974, 1981 and 1995) and in Compendium of Organic Synthetic Methods Vols.1-9 chapter 7 (Wiley 1971-2000).

For the introduction of Si-containing functionalities into thienyl derivatives (step 1) a large variety of synthetic methods are accessible. The chemist skilled in the art will understand that according to the methodology chosen for step 1 different groups T may be

used. Examples of useful T substituents are halogens (such as Cl, Br and I), sulfonates (such as triflates, tosylates and mesylates), phosphates, C₁₋₄ alkyl, C₂₋₄ alkenyl and C₂₋₄ alkynyl.

Manipulation of Si-containing functional groups (step 2) is widely known in the literature. Recent overviews may be found in *The Chemistry of Organosilicon Compounds*, Vols.1-3, S.Patay, Z.Rappaport and Y.Apeloig eds. Wiley, 1989, 1998, 2001 and in *Houben-Weyl Science of Synthesis, Organometallics Vol.4*, I.Fleming ed., G.Thieme 2002.

Examples of manipulations which are especially relevant to the present invention are hydrogenation or reduction of double or triple bonds (or both) in the Si-containing group (see Example 3, step B below), and functional group manipulation on the silicon atom (for example conversion of halogens to alkyl or alkoxy groups).

Literature examples which illustrate some of the methods which may be used with appropriate modification of Step 1 in the above reaction scheme and which are therefore especially relevant to the preparation of compounds of formula (IIa) and (IIb) include *Tetrahedron* 57, 5339 (2001); *Tetrahedron* 56, 2985 (2000); *Tetrahedron* 46, 2632 and 2632 (1990); WO94/12505; *J.Org.Chem.* 64, 2471 (1999); *J.Org.Chem.* 55, 2993 (1990); *J.Org.Chem.* 51, 5286 (1986); US 4777262; *Zh.Obschei Khimi*, 52, 2767 (1982); *Chemica Scripta* 18, 192 (1981); *J.Med.Chem.* 45, 3022 (2002); A.R.M.Allison et Al., *Tetrahedron Letters* 35, 4425 (1994); *Tetrahedron Letters* 25, 81 and 83 (1984); *Tetrahedron Letters* 30, 3735 (1989); *Chemistry Letters* 2001, 386; *Chemistry Letters* 1990, 2175 and references cited therein.

Surprisingly, it has now been found that the novel compounds of formula (I) have, for practical purposes, a very advantageous spectrum of activities for protecting plants against diseases that are caused by fungi as well as by bacteria and viruses.

The compounds of formula (I) can be used in the agricultural sector and related fields of use as active ingredients for controlling plant pests. The novel compounds are distinguished by excellent activity at low rates of application, by being well tolerated by plants and by being environmentally safe. They have very useful curative, preventive and systemic properties and are used for protecting numerous cultivated plants. The compounds of formula (I) can be used to inhibit or destroy the pests that occur on plants or parts of plants (fruit, blossoms, leaves, stems, tubers, roots) of different crops of useful plants, while at the same time protecting also those parts of the plants that grow later, for example, from phytopathogenic microorganisms.

It is also possible to use compounds of formula (I) as dressing agents for the treatment of plant propagation material, in particular of seeds (fruit, tubers, grains) and plant cuttings (for example rice), for the protection against fungal infections as well as against phytopathogenic fungi occurring in the soil.

- 5 Furthermore the compounds according to present invention may be used for controlling fungi in related areas, for example in the protection of technical materials, including wood and wood related technical products, in food storage, in hygiene management, etc.

- The compounds of formula (I) are, for example, effective against the phytopathogenic fungi of the following classes: Fungi imperfecti (for example *Botrytis*, *Pyricularia*,
10 *Helminthosporium*, *Fusarium*, *Septoria*, *Cercospora* and *Alternaria*) and Basidiomycetes (forexample *Rhizoctonia*, *Hemileia*, *Puccinia*). Additionally, they are also effective against the Ascomycetes classes (for example *Venturia* and *Erysiphe*, *Podosphaera*, *Monilinia*, *Uncinula*) and of the Oomycetes classes (for example *Phytophthora*, *Pythium*, *Plasmopara*). Outstanding activity has been observed against powdery mildew (*Erysiphe* spp.). Furthermore, the novel
15 compounds of formula I are effective against phytopathogenic bacteria and viruses (for example against *Xanthomonas* spp, *Pseudomonas* spp, *Erwinia amylovora* as well as against the tobacco mosaic virus).

- Within the scope of present invention, target crops to be protected typically comprise the following species of plants: cereal (wheat, barley, rye, oat, rice, maize, sorghum and related
20 species); beet (sugar beet and fodder beet); pomes, drupes and soft fruit (apples, pears, plums, peaches, almonds, cherries, strawberries, raspberries and blackberries); leguminous plants (beans, lentils, peas, soybeans); oil plants (rape, mustard, poppy, olives, sunflowers, coconut, castor oil plants, cocoa beans, groundnuts); cucumber plants (pumpkins, cucumbers, melons); fibre plants (cotton, flax, hemp, jute); citrus fruit (oranges, lemons, grapefruit, mandarins);
25 vegetables (spinach, lettuce, asparagus, cabbages, carrots, onions, tomatoes, potatoes, paprika); lauraceae (avocado, cinnamomum, camphor) or plants such as tobacco, nuts, coffee, eggplants, sugar cane, tea, pepper, vines, hops, bananas and natural rubber plants, as well as ornamentals.

- The compounds of formula (I) are used in unmodified form or, preferably, together with
30 the adjuvants conventionally employed in the art of formulation. To this end they are conveniently formulated in known manner to emulsifiable concentrates, coatable pastes, directly

sprayable or dilutable solutions, dilute emulsions, wettable powders, soluble powders, dusts, granulates, and also encapsulations, for example, in polymeric substances. As with the type of the compositions, the methods of application, such as spraying, atomising, dusting, scattering, coating or pouring, are chosen in accordance with the intended objectives and the prevailing
5 circumstances. The compositions may also contain further adjuvants such as stabilizers, antifoams, viscosity regulators, binders or tackifiers as well as fertilizers, micronutrient donors or other formulations for obtaining special effects.

Suitable carriers and adjuvants can be solid or liquid and are substances useful in formulation technology, for example, natural or regenerated mineral substances, solvents,
10 dispersants, wetting agents, tackifiers, thickeners, binders or fertilizers. Such carriers are for example described in WO 97/33890.

The compounds of formula (I) are normally used in the form of compositions and can be applied to the crop area or plant to be treated, simultaneously or in succession with further compounds. These further compounds may be, for example, fertilizers or micronutrient
15 donors or other preparations which influence the growth of plants. They may also be selective herbicides as well as insecticides, fungicides, bactericides, nematocides, molluscicides or mixtures of several of these preparations, if desired together with further carriers, surfactants or application promoting adjuvants customarily employed in the art of formulation.

The compounds of formula (I) may be mixed with other fungicides, resulting in some
20 cases in unexpected synergistic activities. Mixing components which are particularly preferred are azoles, such as azaconazole, BAY 14120, bitertanol, bromuconazole, cyproconazole, difenoconazole, diniconazole, epoxiconazole, fenbuconazole, fluquinconazole, flusilazole, flutriafol, hexaconazole, imazalil, imibenconazole, ipconazole, metconazole, myclobutanil, pefurazoate, penconazole, pyrifenox, prochloraz, propiconazole, simeconazole,
25 tebuconazole, tetraconazole, triadimefon, triadimenol, triflumizole, triticonazole; pyrimidinyl carbinole, such as ancymidol, fenarimol, nuarimol; 2-amino-pyrimidines, such as bupirimate, dimethirimol, ethirimol; morpholines, such as dodemorph, fenpropidine, fenpropimorph, spiroxamine, tridemorph; anilinopyrimidines, such as cyprodinil, mepanipyrim, pyrimethanil; pyrroles, such as fenpiclonil, fludioxonil; phenylamides, such as benalaxyl, furalaxyl, meta-
30 laxyl, R-metalaxyl, ofurace, oxadixyl; benzimidazoles, such as benomyl, carbendazim, debacarb, fuberidazole, thiabendazole; dicarboximides, such as chlozolate, dichlozoline,

iprodione, myclozoline, procymidone, vinclozoline; carboxamides, such as carboxin, fenfuram, flutolanil, mepronil, oxycarboxin, thifluzamide; guanidines, such as guazatine, dodine, iminoctadine; strobilurines, such as azoxystrobin, kresoxim-methyl, metominostrobin, SSF-129, trifloxystrobin, picoxystrobin, BAS 500F (proposed name pyraclostrobin), BAS 520; dithiocarbamates, such as ferbam, mancozeb, maneb, metiram, propineb, thiram, zineb, ziram; N-halomethylthiotetrahydrophthalimides, such as captafol, captan, dichlofluanid, fluoromides, folpet, tolyfluanid; Cu-compounds, such as Bordeaux mixture, copper hydroxide, copper oxychloride, copper sulfate, cuprous oxide, mancopper, oxine-copper; nitrophenol-derivatives, such as dinocap, nitrothal-isopropyl; organo-p-derivatives, such as edifenphos, iprobenphos, isoprothiolane, phosdiphen, pyrazophos, tolclofos-methyl; various others, such as acibenzolar-S-methyl, anilazine, benthiavalicarb, blasticidin-S, chinomethionate, chloroneb, chlorothalonil, cyflufenamid, cymoxanil, dichlone, diclomezine, dicloran, diethofencarb, dimethomorph, SYP-LI90 (proposed name: flumorph), dithianon, ethaboxam, etridiazole, famoxadone, fenamidone, fenoxanil, fentin, ferimzone, fluazinam, flusulfamide, fenhexamid, fosetyl-aluminium, hymexazol, iprovalicarb, IKF-916 (cyazofamid), kasugamycin, methasulfocarb, metrafenone, nicobifen, pencycuron, phthalide, polyoxins, probenazole, propamocarb, pyroquilon, quinoxifen, quintozone, sulfur, triazoxide, tricyclazole, triforine, validamycin, zoxamide (RH7281).

A preferred method of applying a compound of formula (I), or an agrochemical composition which contains at least one of said compounds, is foliar application. The frequency of application and the rate of application will depend on the risk of infestation by the corresponding pathogen. However, the compounds of formula I can also penetrate the plant through the roots via the soil (systemic action) by drenching the locus of the plant with a liquid formulation, or by applying the compounds in solid form to the soil, for example, in granular form (soil application). In crops of water rice such granulates can be applied to the flooded rice field. The compounds of formula I may also be applied to seeds (coating) by impregnating the seeds or tubers either with a liquid formulation of the fungicide or coating them with a solid formulation.

A formulation [that is, a composition containing the compound of formula (I)] and, if desired, a solid or liquid adjuvant, is prepared in a known manner, typically by intimately

mixing and/or grinding the compound with extenders, for example solvents, solid carriers and, optionally, surface active compounds (surfactants).

The agrochemical formulations will usually contain from 0.1 to 99% by weight, preferably from 0.1 to 95% by weight, of the compound of formula I, 99.9 to 1% by weight, preferably 99.8 to 5% by weight, of a solid or liquid adjuvant, and from 0 to 25% by weight, preferably from 0.1 to 25% by weight, of a surfactant.

Advantageous rates of application are normally from 5g to 2kg of active ingredient (a.i.) per hectare (ha), preferably from 10g to 1kg a.i./ha, most preferably from 20g to 600g a.i./ha. When used as seed drenching agent, convenient dosages are from 10mg to 1g of active substance per kg of seeds.

Whereas it is preferred to formulate commercial products as concentrates, the end user will normally use dilute formulations.

The following non-limiting Examples illustrate the above-described invention in more detail.

15 EXAMPLE 1

This Example illustrates the preparation of Compound Numbers 27.2 and 1.14.

Step A

4-Bromo-3-thiophene carboxylic acid (prepared according to J.Heterocyclic Chem. 1999,36,761) (2g), triethylamine (1.5g) and acetone (15 ml) were cooled to -10°C and ethylchloroformiate (1.15g) was added at this temperature. The reaction mixture was stirred at room temperature for 2 hours, cooled to 0°C and sodium azide (0.97g) dissolved in water (5ml) was added slowly. After 2 hours at room temperature the mixture was filtered, the acetone was evaporated and the residue was diluted with dichloromethane, washed with water and dried over sodium sulfate. Evaporation of the solvent yielded 1.7g of an off-white powder which was directly used in the next step

Step B: 4-Bromo-3-thiophene-formamide

Formic acid (30ml) was heated to reflux and the product of Step A, dissolved in formic acid (10ml), was added dropwise. The mixture was heated for 30 minutes until gas evolution stopped. The formic acid was removed under vacuum and then the residue was diluted with ethyl acetate and washed with saturated sodium bicarbonate solution. After drying and

removal of the solvent the product was chromatographed on silica gel (hexane:ethylacetate 2:1) to yield 1.1g of product (m.p.108-110°C).

Step C: 4-(Trimethylsilyl)ethinyl-3-thiophene formamide

The product from step B (1g), tributylstannyl(trimethylsilyl)acetylene (2.16g) and
5 palladium(tetrakis)triphenylphosphine (0.28g) were placed into degassed toluene (20ml) and the mixture was heated under reflux for 16 hours under an atmosphere of nitrogen. After cooling, the solvent was removed and the residue was chromatographed on silica gel (hexane:ethylacetate 3:1). After three consecutive crystallisations from hexane, 0.3g of 4-(trimethylsilyl)ethinyl-3-thiophene formamide was obtained (m.p.90-94°C).

10 Step D : Compound Number 27.2

The product obtained from step C was hydrogenated in THF over Pd on charcoal (10%) (0.3g) until hydrogen uptake stopped. Evaporation of the solvent yielded Compound Number 27.2 (0.28g).

Step E: Compound Number 1.14

15 Sodium hydride (0.06g) was dispersed in THF (10ml) and the mixture was cooled with an ice bath. Compound Number 27.2 (0.14g), dissolved in THF (5ml), was added and the suspension was stirred at 0-5°C for 30 minutes. N-Methyl-3-trifluoromethyl-4-chlorocarbonyl-pyrazole (0.2g) was added at this temperature and the reaction mixture was stirred at room temperature overnight. The reaction mixture was poured on to water and was
20 extracted with ethylacetate. The organic phase was washed with water, dried over sodium sulfate and the solvent was evaporated. The residue was treated with an excess of sodium methylate in methanol and then, after 15 minutes at room temperature, the mixture was acidified with HCl and extracted with ethyl acetate. After washing with saturated sodium bicarbonate, drying over sodium sulfate and removing the solvent, the residue was
25 chromatographed on silica gel (hexane:ethyl acetate 1:1) and recrystallised from hexane, yielding Compound Number 1.14 (0.072g).

EXAMPLE 2

This Example illustrates the preparation of Compound Numbers 28.2, 2.85 and 2.16.

Step A: 2-Iodo-3-formylaminothiophene

30 3-Formylamino-thiophene (21g), N-iodosuccinimide (37.2g) and tetrachlormethane (200ml) were refluxed together for 3 hours. The reaction mixture was cooled to 0-5°C and

filtered. The solids were dissolved in ethyl acetate, washed three times with sodium hydroxide (10% solution), washed with brine and dried over sodium sulfate. Evaporation of the solvents and chromatography of the residue over silica gel (hexane:ethyl acetate 2:1) yielded 37.8g of product (m.p.100-107°C) which was directly used in Step B.

5 Step B: 2-(Trimethylsilylethynyl)-3-formamido-thiophene.

The product from step A (37.8g) was dissolved in a mixture of DMF (500ml) and triethylamine (150ml). CuI (1.4g) and bis(triphenylphosphino)palladiumdichloride (5.2g) were added under nitrogen and after 15 minutes ethynyl trimethylsilane (22g) was added dropwise. After 6 hours the solvents were removed under vacuum and the residue was
10 chromatographed on silica gel (hexane:ethyl acetate 2:1) to yield 2-(trimethylsilylethynyl)-3-formamido-thiophene (31.78g) (m.p.87-93°C).

Step C: Compound Number 28.2

Hydrogenation of the product from Step B (31.78g) in THF over Pd (30g) on charcoal, as described for Compound Number 27.2 in Step D of Example 1, yielded Compound
15 Number 28.2 (26.3g).

Step D: Compound Numbers 2.85 and 2.16

Compound 28.2 (23.4g) was treated with sodium hydride (7.4g) and N-methyl-3-difluoromethyl-4-chlorocarbonyl pyrazole (30g) as described in Step E of Example 1, to form a reaction mixture.

20 In order to prepare Compound Number 2.85, one tenth of the reaction mixture was poured on to water and was extracted with ethylacetate. The organic phase was washed with water, dried over sodium sulfate and the solvent was evaporated. The residue was chromatographed on silica gel (hexane:ethylacetate 1:1) and Compound Number 2.85 was isolated by removal of the solvent; yield: 0.6g.

25 In order to prepare Compound Number 2.16, the reaction mixture was treated with sodium hydroxide (100ml of a 10% w/w solution in water) and stirred over night. The reaction mixture was diluted with saturated ammonium chloride solution, extracted with ethyl acetate, dried over sodium sulfate and the solvent was removed under reduced pressure. The residue was recrystallised with hexane:toluene (3:1) to yield Compound Number 2.16 (27.7g).

EXAMPLE 3

This Example illustrates the preparation of Compound Number 6.1.

Step A

3- Formylaminothiophene (Bull.Soc.Chim.Fr. 1976,151) (2g), dissolved in dry THF
5 (20ml), was dropped into a freshly prepared solution of lithium diisopropylamine (0.035mol)
in THF (35ml) at -70°C. The solution was warmed to -20°C and stirred for 30 minutes, then
cooled again to -70°C, whereupon trimethylsilylchloride (5ml) was added. After stirring for
an additional hour at this temperature the mixture was slowly warmed to room temperature
and stirred for 4 hours. Quenching with a saturated solution of ammonium chloride,
10 extracting with ethyl acetate, drying over sodium sulfate, removing the solvent and applying
chromatography on silica gel (hexane:ethyl acetate 2:1) yielded 2-trimethylsilyl-3-
formylaminothiophene (1.2g) (m.p.74-76°C).

Step B: Compound Number 6.1

Preparation of Compound Number 6.1 was carried out according to Step D of Example
15 2 using the above intermediate from Step A (0.4g), sodium hydride (0.13g), 2-methyl-4-
trifluormethyl-5-chlorocarbonyl-thiazole (0.69g) and sodium hydroxide solution (1.5 ml of a
2 molar solution). Chromatography on silica gel (hexane:ethyl acetate 4:1) yielded
Compound Number 6.1 (0.29g).

FORMULATION EXAMPLES FOR COMPOUNDS OF FORMULA (I)

20 Working procedures for preparing formulations of the compounds of formula I such as
Emulsifiable concentrates, Solutions, Granulates, Dusts and Wettable powders are described
in WO 97/33890.

BIOLOGICAL EXAMPLES: FUNGICIDAL ACTIONS

Example B-1: Action against *Puccinia recondita* / wheat (Brownrust on wheat)

25 1 week old wheat plants cv. Arina were treated with the formulated test compound
(0.02% active ingredient) in a spray chamber. One day after application wheat plants were
inoculated by spraying a spore suspension (1×10^5 uredospores/ml) on the test plants. After an
incubation period of 2 days at 20°C and 95% r. h. plants were kept in a greenhouse for 8 days
at 20°C and 60% r.h. The disease incidence was assessed 10days after inoculation.

30 Infestation was prevented virtually completely (0-5% infestation) with each of
Compounds 1.14, 1.16, 2.1, 2.2, 2.14, 2.16, 2.75, 2.76, 2.84, 2.85 and 6.01.

Example B-2: Action against *Podosphaera leucotricha* / apple (Powdery mildew on apple)

5 week old apple seedlings cv. McIntosh were treated with the formulated test compound (0.002% active ingredient) in a spray chamber. One day after application apple plants were inoculated by shaking plants infected with apple powdery mildew above the test plants. After an incubation period of 12 days at 22°C and 60% r.h. under a light regime of 14/10 hours (light/dark) the disease incidence was assessed.

Compounds 2.14 and 2.16 each exhibited strong efficacy (<20% infestation).

Example B-3: Action against *Venturia inaequalis* / apple (Scab on apple)

4 week old apple seedlings cv. McIntosh were treated with the formulated test compound (0.02% active ingredient) in a spray chamber. One day after application apple plants were inoculated by spraying a spore suspension (4×10^5 conidia/ml) on the test plants. After an incubation period of 4 days at 21°C and 95% r.h. the plants were placed for 4 days at 21°C and 60% r.h. in a greenhouse. After another 4 day incubation period at 21°C and 95% r.h. the disease incidence was assessed.

Compounds 2.14 and 2.16 each exhibited strong efficacy (<20% infestation).

Example B-4: Action against *Erysiphe graminis* / barley (Powdery mildew on barley)

1 week old barley plants cv. Express were treated with the formulated test compound (0.02% active ingredient) in a spray chamber. One day after application barley plants were inoculated by shaking powdery mildew infected plants above the test plants. After an incubation period of 6 days at 20°C / 18°C (day/night) and 60% r.h. in a greenhouse the disease incidence was assessed.

Compounds 1.14, 1.16, 2.14, 2.16, 2.75, 2.76 and 2.85 each exhibit strong efficacy (<20% infestation).

Example B-5: Action against *Botrytis cinerea* / tomato (Botrytis on tomatoes)

4 week old tomato plants cv. Roter Gnom were treated with the formulated test compound (0.02% active ingredient) in a spray chamber. Two days after application tomato plants were inoculated by spraying a spore suspension (1×10^5 conidia/ml) on the test plants. After an incubation period of 4 days at 20°C and 95% r.h. in a growth chamber the disease incidence was assessed.

Compounds 1.14, 1.16, 2.14, 2.16, 2.75, 2.76 and 2.85 each exhibit strong efficacy (<20% disease incidence).


Example B-6 Action against *Septoria nodorum* / wheat (Septoria leaf spot on wheat)

1 week old wheat plants cv. Arina were treated with the formulated test compound (0.02% active ingredient) in a spray chamber. One day after application wheat plants were inoculated by spraying a spore suspension (5×10^5 conidia/ml) on the test plants. After an
5 incubation period of 1 day at 20°C and 95% r.h. plants are kept for 10 days at 20°C and 60% r.h. in a greenhouse. The disease incidence was assessed 11 days after inoculation.

Compounds 1.14, 1.16, 2.14, 2.16, 2.75, 2.76 and 2.85 each show good activity in this test (<60% disease incidence).

CLAIMS

1. A compound of formula (I)



- 5 where Het is a 5- or 6-membered heterocyclic ring containing one to three heteroatoms, each independently selected from oxygen, nitrogen and sulphur, the ring being substituted by one to three groups R^4 ; R^1 is hydrogen, optionally substituted (C₁₋₄)alkyl, formyl, optionally substituted (C₁₋₄)alkylC(=O), optionally substituted (C₁₋₄)alkylC(=O)O, optionally substituted (C₁₋₄)alkoxy(C₁₋₄)alkyl, optionally substituted
- 10 allyl, optionally substituted propargyl or optionally substituted allenyl; each R^2 is, independently, halogen, optionally substituted (C₁₋₄)alkyl, optionally substituted (C₁₋₄)alkoxy or optionally substituted (C₁₋₄)alkoxy(C₁₋₄)alkyl; R^3 is either at position 2 or at position 4 of the thiophene ring and is an organic group containing three to thirteen carbon atoms and at least one silicon atom and, optionally, one to three heteroatoms,
- 15 each independently selected from oxygen, nitrogen and sulphur, and is optionally substituted by one to four independently selected halogen atoms; each R^4 is, independently, halogen, C₁₋₃ alkyl, C₁₋₃ haloalkyl, C₁₋₃alkoxy(C₁₋₃)alkyl or cyano; r is 0, 1 or 2; and X is O or S; or an N-oxide thereof.
- 20 2. A compound of formula (I) as defined in claim 1 provided that the Het ring is not 1,2,3-triazole when X is O.
3. A compound of formula (I) as defined in claim 1 or 2 where R^1 is hydrogen, propargyl, allenyl, formyl, CH₃C(=O), C₂H₅C(=O) or CH₃OCH₂C(=O).
- 25 4. A compound of formula (I) as defined in claim 1, 2 or 3 where each R^2 is independently selected from halogen, methyl, trifluoromethyl and trifluoromethoxy.

5. A compound of formula (I) as defined in claim 1, 2, 3 or 4 where Het is pyrazolyl, pyrrolyl, thiophenyl, furyl, thiazolyl, isothiazolyl, oxazolyl, isoxazolyl, triazolyl, pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, 5,6-dihydropyranyl or 5,6-dihydro-1,4-oxathiinyl each being substituted by one to three groups R^4 .


5

6. A compound of formula (I) as defined in claim 1, 2, 3, 4 or 5 where R^3 is an aliphatic, saturated or unsaturated group containing three to thirteen carbon atoms and at least one silicon atom and, optionally, one to three heteroatoms, each independently selected from oxygen, nitrogen and sulphur, and is optionally substituted by one to four independently selected halogen atoms.


10

7. A compound of formula (I) as defined in claim 1, 2, 3, 4, 5 or 6 where X is oxygen.

8. A compound of formula (IIa) or (IIb)



(IIa)



(IIb)

15

where R^3 is SiMe_2Et , $\text{SiMe}_2\text{CHMe}_2$, $\text{SiMe}_2\text{CH}_2\text{CHMe}_2$, $\text{SiMe}_2\text{CH}_2\text{CMe}_3$, $\text{CH}=\text{CHSiMe}_3$, $(\text{CH}_2)_2\text{SiMe}_3$, $(\text{CH}_2)_2\text{SiMe}_2\text{Et}$, $(\text{CH}_2)_2\text{SiMe}_2\text{CHMe}_2$, $(\text{CH}_2)_2\text{SiMe}_2\text{CMe}_3$, $(\text{CH}_2)_2\text{SiMe}_2\text{CH}_2\text{CHMe}_2$, $\text{CHMeCH}_2\text{SiMe}_3$, $\text{CHMeCH}_2\text{SiMe}_2\text{Et}$, $\text{CH}_2\text{CHMeSiMe}_3$, $\text{CMe}_2\text{CH}_2\text{SiMe}_3$, $(\text{CH}_2)_3\text{SiMe}_3$; and A is NH_2 , $\text{NHCH}(\text{O})$, optionally substituted $(\text{C}_{1-4})\text{alkylC}(=\text{O})\text{NH}$, optionally substituted $(\text{C}_{1-4})\text{alkylOC}(=\text{O})\text{NH}$, halogen or $\text{N}=\text{C}(\text{C}_6\text{H}_5)_2$; provided that A is not halogen when R^3 is SiMe_3 .

20

9. A composition for controlling microorganisms and preventing attack and infestation of plants therewith, wherein the active ingredient is a compound of formula (I) as claimed in claim 1 together with a suitable carrier.

25

10. A method of controlling or preventing infestation of cultivated plants by phytopathogenic microorganisms by application of a compound of formula (I) as claimed in claim 1 or a composition as claimed in claim 9 to plants, to parts thereof or the locus thereof.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP2004/004194

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D409/12 C07D417/12 C07D413/12 C07D411/12 C07D333/36
C07D333/28 A01N43/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 96/18631 A (MONSANTO CO ; WONG SAI CHI (US); PHILLION DENNIS PAUL (US); SHORTT BAR) 20 June 1996 (1996-06-20) page 1, paragraph 1 claim 1	1,9
A	WO 99/62915 A (MONSANTO CO) 9 December 1999 (1999-12-09) page 1, line 9 - line 12 claim 1	1,9
A,P	WO 2004/018438 A (EHRENFREUND JOSEF ; TOBLER HANS (CH); WALTER HARALD (CH); SYNGENTA PAR) 4 March 2004 (2004-03-04) claim 1	1,9

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

19 August 2004

Date of mailing of the international search report

30/08/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Fanni, S

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP2004/004194

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9618631	A	20-06-1996	US 5486621 A	23-01-1996
			AP 725 A	29-01-1999
			AT 192750 T	15-05-2000
			AU 688639 B2	12-03-1998
			AU 4108096 A	03-07-1996
			BG 62853 B1	29-09-2000
			BG 101774 A	30-10-1998
			BR 9510029 A	02-06-1998
			CA 2207749 A1	20-06-1996
			CN 1176642 A , B	18-03-1998
			CZ 9701826 A3	17-12-1997
			DE 69516899 D1	15-06-2000
			DE 69516899 T2	21-12-2000
			DK 797578 T3	14-08-2000
			EE 9700126 A	15-12-1997
			EP 0797578 A1	01-10-1997
			ES 2148582 T3	16-10-2000
			FI 972524 A	13-08-1997
			GR 3033935 T3	30-11-2000
			HK 1009137 A1	24-11-2000
			HR 950597 A1	31-10-1997
			HU 77794 A2	28-08-1998
			JP 11510472 T	14-09-1999
			MD 970248 A	30-09-1999
			NO 972730 A	15-08-1997
			NZ 296458 A	29-06-1999
			PL 320735 A1	27-10-1997
			PT 797578 T	31-10-2000
			RO 117795 B1	30-07-2002
			RU 2145962 C1	27-02-2000
			SI 797578 T1	31-08-2000
			SK 75797 A3	14-01-1998
			TJ 278 B	13-12-2000
			WO 9618631 A1	20-06-1996
			ZA 9510665 A	16-07-1996
WO 9962915	A	09-12-1999	AT 223422 T	15-09-2002
			AU 762985 B2	10-07-2003
			AU 4418399 A	20-12-1999
			BR 9911624 A	30-10-2001
			CA 2334221 A1	09-12-1999
			CN 1314907 T	26-09-2001
			DK 1084130 T3	21-10-2002
			EP 1084130 A1	21-03-2001
			ES 2183568 T3	16-03-2003
			PL 345161 A1	03-12-2001
			PT 1084130 T	31-01-2003
			RU 2220973 C2	10-01-2004
			WO 9962915 A1	09-12-1999
			US 6140511 A	31-10-2000
			US 6291513 B1	18-09-2001
WO 2004018438	A	04-03-2004	WO 2004018438 A2	04-03-2004